

AD _____
(Leave blank)

Award Number: W81XWH-07-1-0623

TITLE: A genetic model for the breast cancer microenvironment

PRINCIPAL INVESTIGATOR: James F. Amatruda, M.D., Ph.D.

CONTRACTING ORGANIZATION: University of Texas Southwestern
Medical Center at Dallas
Dallas, TX 75390

REPORT DATE: September 2008

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: (Check one)

☒ Approved for public release; distribution unlimited

☐ Distribution limited to U.S. Government agencies only;
report contains proprietary information

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

Table of Contents

	<u>Page</u>
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusion.....	7
References.....	8
Appendices.....	9


INTRODUCTION

Despite improved treatments, metastatic breast cancer kills more than 40,000 women each year in the US. Analysis of metastatic breast cancer cell lines has revealed many factors produced by the tumor cells that promote metastasis, for example matrix metalloproteinases, adhesion molecules and angiogenic growth factors. In contrast, relatively little is known about what factors in the host contribute to the establishment of metastases. Recent intriguing evidence points to novel mechanisms, such as α v β 3 integrin expression on endothelial cells, expression of chemokine CXCL12 by target tissues and p53 mutations in tumor-associated stroma, as contributors to invasiveness and metastatic potential (Kurose, Gilley et al. 2002; Dewan, Ahmed et al. 2006; Gupta and Massague 2006). It is critically important to discover other such host factors; however the tumor microenvironment has been difficult to study because it cannot be manipulated as easily as can the cell lines. To address this problem we have turned to the zebrafish, a powerful, genetically tractable vertebrate model system with cancer biology very similar to human (Amatruda, Shepard et al. 2002). The goal of this study is to understand how the host microenvironment influences the development of breast cancer metastasis.


We hypothesize that, when placed in zebrafish embryos as xenografts, human breast cancer cells will recruit host factors to survive, proliferate and spread. Recently, it was shown that human melanoma cell xenografts survive and stimulate angiogenesis when placed in zebrafish embryos (Haldi, Ton et al. 2006). The overall approach of this project is to use knockdown and mutagenesis techniques in zebrafish embryos to systematically alter the tumor microenvironment, assessing the impact on growth and invasiveness of the xenografted human breast cancer cells.

BODY:

In the first phase of this project, we chose to focus on the role of chemokine signaling in the interaction of breast cancer cells with their environment. The chemokine receptor CXCR4b is expressed in breast cancer cells and is associated with tumor growth, spread and angiogenesis (Dewan, Ahmed et al. 2006; Liang, Brooks et al. 2007). The ligand for CXCR4b is known as CXCL12 or Sdf-1a (for Stromal-derived factor-1a). Our goal was to create an experimental system whereby we could investigate the role of Sdf-1a expression in the microenvironment. Specifically, we set up a system to manipulate Sdf-1a expression in zebrafish embryos. The experimental scheme is shown in **Figure 1**. Using a robust antisense technology known as a morpholino, we can knock down endogenous *SDF-1* expression; this approach leads to stable knockdown for 4-5 days. Separately, we inject an embryo with mRNA encoding zebrafish Sdf-1a along with a fluorescent label. We then transplant labeled cells expressing high levels of *SDF-1* into the knockdown embryos. This creates a localized area of *SDF-1* expression. The purpose of this proof-of-principle experiment is to assess whether misexpression of Sdf-1a is sufficient to alter the migration of Primordial Germ Cells (PGCs). To more easily visualize PGCs in living animals, we take advantage of the *askopos*:GFP transgenic line, in which the PGCs are tagged with GFP. We then assess whether PGCs (green) migrate toward the *SDF-1*-expressing cells (red).



To demonstrate successful knockdown, we injected *Sdf-1a* morpholino into *askopos:GFP* transgenics and assessed PGC migration. **Figure 2** shows that knockdown of *Sdf-1a* leads to aberrant migration of PGCs.



We then proceeded to carry out the experimental scheme outlined in Figure 1. **Figure 3** shows an embryo in which the GFP-labeled PGCs have ectopically migrated to sites of high Sdf-1a expression, as identified by Texas Red dextran-labeled transplanted cells.





Figure 3. PGCs migrate ectopically toward misexpressed Sdf-1a. PGCs (green) in an *askopos*-GFP *SDF-1* knockdown embryo migrate toward exogenous cells (red) expressing high levels of *SDF-1*.

Methods: 1-cell stage *askopos*-GFP transgenic zebrafish embryos were injected with an *SDF-1* morpholino. 1-cell stage wild-type zebrafish embryos were injected with *SDF-1* mRNA and Texas Red Dextran. The *SDF-1* positive cells were transplanted into the *askopos*-GFP *SDF-1* knockdown embryo at 3 hours post-fertilization, creating localized *SDF-1* expression. Cells were visualized 24 hours post-fertilization with epifluorescence microscopy.

A key aspect of the experimental scheme is showing that we can transplant human tumor xenografts into zebrafish embryos, and the cells will survive and be visible when fluorescently labeled. **Figure 4** demonstrates labeled human tumor xenografts in live zebrafish embryos. We also went on to characterize CXCR4b and Sdf-1a expression in a large collection of primary breast cancer cell lines at UT Southwestern Medical Center. Currently, we are growing these cells and attempting to achieve stable, fluorescently-labeled cell lines from each. Once the lines are obtained, we will carry out the experiment to observe the effect of Sdf-1a misexpression on human breast cancer cell proliferation and migration (experimental schema shown in **Figure 5**.)

KEY RESEARCH ACCOMPLISHMENTS

- We have generated a system whereby we can selectively alter a key component of the tumor microenvironment (Sdf-1a expression)
- We have shown that misexpressed Sdf-1a in the embryo microenvironment serves as a guidance cue for PGC migration, which depends on CXCR4b/Sdf-1a interaction.
- We have demonstrated that labeled human tumor cells can be xenografted into zebrafish embryos
- We have identified breast cancer cells lines with varying expression of CXCR4b and Sdf-1a expression levels
-



REPORTABLE OUTCOMES:


Abstract submitted to 2008 Era of Hope Meeting, Baltimore, MD

Poster Presentation 2008 Era of Hope Meeting, Baltimore, MD

CONCLUSION:

We have taken the important initial steps into creating a genetically-tractable model of the tumor microenvironment, focusing on the critical CXCR4b/Sdf-1a interaction. Importantly, we were able to use endogenous Primordial Germ Cell Migration as a reporter to show that we successfully modified Sdf-1a expression in a way that reveals CXCR4b-mediated migratory events. We now need to repeat the experiment using labeled human breast cancer cells as the reporter. Once successfully established, this model could be used as a rapid, convenient and physiological model of CXCR4b-dependent breast cancer cell migration. The main

purpose of such a model would be to test molecular hypotheses (e.g. by knocking down key target genes in breast cancer cells, or by knocking down or misexpressing genes in the embryo microenvironment) or for relatively high-throughput screening or novel agents that could inhibit breast cancer cell migration and therefore potentially abrogate metastasis.



REFERENCES:

- Amatruda, J. F., J. L. Shepard, et al. (2002). "Zebrafish as a cancer model system." Cancer Cell 1(3): 229-31.
- Dewan, M. Z., S. Ahmed, et al. (2006). "Stromal cell-derived factor-1 and CXCR4 receptor interaction in tumor growth and metastasis of breast cancer." Biomed Pharmacother 60(6): 273-6.
- Gupta, G. P. and J. Massague (2006). "Cancer metastasis: building a framework." Cell 127(4): 679-95.
- Haldi, M., C. Ton, et al. (2006). "Human melanoma cells transplanted into zebrafish proliferate, migrate, produce melanin, form masses and stimulate angiogenesis in zebrafish." Angiogenesis 9(3): 139-51.
- Kurose, K., K. Gilley, et al. (2002). "Frequent somatic mutations in PTEN and TP53 are mutually exclusive in the stroma of breast carcinomas." Nat Genet 32(3): 355-7.
- Liang, Z., J. Brooks, et al. (2007). "CXCR4/CXCL12 axis promotes VEGF-mediated tumor angiogenesis through Akt signaling pathway." Biochem Biophys Res Commun 359(3): 716-22.

Appendices:

None